

Monitoring NAD(P)H levels and their relevance to bioenergetics using Fluorescence-Based Strategies in salivary samples: a possible biomarker in the detection of cancer and precancer

Pallotta ML EUSTIM 2016

Pallotta ML L-proline homeostasis, substrate channeling and mitochondrial flavine turnover in microenvironment: phenotypic landscape of *Saccharomyces cerevisiae* for pharmacological screening *in silico/in vitro* Abstract submitted and approved to 41st Febs Congress 2016, Molecular and Systems Biology for a Better life, Area Epigenetics and Cancer 3-8 September 2016 Ephesus / Kuşadası Turkey

Rossi F, **Pallotta ML** Bacteriocin Producing Cultures: A Sustainable Way for Food Safety Improvement and Probiotics with Additional Health Promoting Effects

International Journal of Medical and Biological Frontiers (2016) 22 (1):59-91, ISSN 1081-3829

Pallotta ML 68 genes for Yeast Flavoproteoma: updates of Flavin Biosynthesis, Transport and Catabolism in *Saccharomyces cerevisiae* mitochondria^{3rd} International Conference on Integrative Biology, August 04-06 Valence Spain 2015, Biol Syst Open Access ISSN: 2329-6577doi: [10.4172/2329-6577.S1.003](https://doi.org/10.4172/2329-6577.S1.003)

Pallotta ML The Aquarian Age in cancer therapy: biobankers, biomarkers, and much more ↔ getting of the root of the problem ↔ NAD⁺ an ancient metabolite with surprising modernity ↔ Mitochondrial NAD pool, tumour microenvironment and metastasis

Cell Symposia, Multifaceted Mitochondria, Chicago July 19-21, 2015.

Pallotta ML Boosting NAD(P)⁺ biosynthesis with NAD(P)⁺ intermediates and monitoring mitochondrial NAD(P)⁺/NAD(P)H pool by means of Fluorescence-Based techniques could be a strategy for preventing and treating Woman's Cancers. FEBS 2015 Berlin July 4-9, 2015 Accession Number: WOS:000362570605033.

Rossi F., Colavita G., Amadoro C. and **Pallotta ML** Traditional Italian dairy products: a flavourful source of naturally occurring bacteria with beneficial effects on health. Bageco 2015, 13th Symposium on Bacterial Genetics and Ecology, The Microbiol Continuity Across Evolving Ecosystems, 14-18 June Milan.

Pallotta ML NMN, an intermediate in the salvage pathway, helps to increase the intramitochondrial NAD⁺ concentration in yeast and influences NAD⁺/NADH ratio: possible function as metabolic read-out 65. Mosbacher Kolloquium - Molecular Protein Quality Control in Health, Aging and Disease, German Society for Biochemistry and Molecular Biology, Mosbach Germany 26th-29th March 2014 DOI: 10.3288/contoo.paper.2990

Pallotta ML Exogenous NMN fuels mitochondrial NAD pool in *Saccharomyces cerevisiae*: probably role as sensitive and dynamic regulator of energy metabolism in response to a cellular microenvironment and its application in human health

Giornata scientifica del Gruppo Membrane e Bioenergetica della SIBBM Matera 28-29 Maggio 2014

Pallotta ML 68 genes for yeast flavoproteoma: general aspects in Flavin biosynthesis, transport and catabolism in *Saccharomyces cerevisiae* mitochondria

2nd International Symposium on Energy challenges and Mechanics- Session Bioenergetics

University of Aberdeen, Scotland, United kingdom

Pallotta ML Mitochondrial NAD pool ad metastasis
Cells, proteins and bioprocessing (Nature) London (England) 17th-19th June 2014

Pallotta ML L-Proline uptake in *Saccharomyces cerevisiae* mitochondria can contribute to bioenergetics during nutrient stress as alternative mitochondrial fuel
World Journal of Microbiology and Biotechnology 30(1):19-21 DOI 10.1007/s11274-013-1415-0

Pallotta ML New insights into L-Proline uptake and mitochondrial energetic metabolism in *Saccharomyces cerevisiae*
26th International Conference on Yeast Genetics and Molecular Biology. Frankfurt/Main, Germany. August 29-September 3, 2013.
Yeast (Chichester, England) Volume: 30 Suppl 1 Pages: S22-253 Published: 2013-Sep

Pallotta ML Mitochondrial involvement to methylglyoxal detoxification: D-Lactate/Malate antiporter in *Saccharomyces cerevisiae*
Antonie van Leeuwenhoek (2012) 102:163–175. DOI 10.1007/s10482-012-9724-0

Pallotta ML Evidence for the presence of a FAD pyrophosphatase and a FMN phosphohydrolase in yeast mitochondria: a possible role in flavin homeostasis
Yeast. 2011 28(10): 693-705. doi: 10.1002/yea.1897

C. Di Martino, **ML Pallotta** Mitochondria-localized NAD biosynthesis by nicotinamide mononucleotide adenyltransferase in Jerusalem artichoke (*Helianthus tuberosus* L.) heterotrophic tissues
Planta 2011 234(4): 657-70 doi:10.1007/s00425-011-1428-6

Pallotta ML *Saccharomyces cerevisiae* mitochondria contain enzymes capable of hydrolyzing FMN and FAD to riboflavin: probably function in flavoprotein deflavination and reconstitution in cell grown under glucose limitation.
35th FEBS Congress, Gothenburg, Sweden 2010
FEBS Journal (2010) Vol: 277: 221-221

ML Pallotta and C Di Martino A new mitochondrial three-component pyridine nucleotide pathway, namely a NMN→NAD→NADP route, is functional in plant heterotrophic tissues
35th FEBS Congress, Gothenburg, Sweden 2010.
FEBS Journal (2010) Vol: 277: 221-221

C Di Martino, **ML Pallotta** and G Palumbo Effect of phosphorus availability on nitrogen assimilation in durum wheat mycorrhized plants.
2nd Congress of the Italian Society of Plant Biology SIBV, Rome 2010

Pallotta ML Plant cells respond to ageing by implementing an emergency survival strategy: increase mitochondrial pyridine dinucleotides pool via NMN-AT.

MLP

Third SMBBM International Congress of Biochemistry, IUBMB Special Meeting on Plant Stress & 6th Congress of FASBMB 2009

Pallotta ML *Saccharomyces cerevisiae* as a model system for studying mitochondria natural flavin catabolism.

16th International Symposium on Flavins and Flavoproteins June 2008

ML. Pallotta and C. Di Martino Existence of intramitochondrial nicotinamide mononucleotide adenyl-transferase activity, which allows for NAD⁺ synthesis from NMN and endogenous ATP in aged-dehydrated slices tubers of *Helianthus tuberosus*.

3rd Cell Stress Society International Congress on Stress responses in Biology and Medicine and 2nd Word Conference of Stress 23-26 August 2007 Budapest (Hungary) Cell Stress & Chaperones Online ISSN: 1466-1268 Print ISSN: 1355-8145 Vol. 12, Issue 2 (Summer 2007).

ML. Pallotta, C. Di Martino Mitochondrion participates in the pathway of NAD(P) biosynthesis in *Saccharomyces cerevisiae*.

Federazione Italiana Scienze della Vita, IX National Congress September 2007

C. Di Martino, **ML Pallotta**

New insights into the NAD synthesis in mitochondria isolated from fresh and aged-dehydrated tubers of Jerusalem artichoke

Federazione Italiana Scienze della Vita, IX National Congress September 2007

ML Pallotta, C Di Martino Age-related changes in proline mitochondrial transport and metabolism in Jerusalem artichoke tubers

Federazione Italiana Scienze della Vita, IX National Congress September 2007

C. Di Martino, R. Pizzuto, **ML. Pallotta**, A. De Santis and S. Passarelli Mitochondrial transport in proline catabolism in plants: the existence of two separate translocators in mitochondria isolated from durum wheat seedlings

Planta (2006) 223(6):1123-1133.

Pallotta ML D-Lactate/malate antiporter, in response to the cellular carbonyl stress mediated by methylglyoxal, in yeast mitochondria.

FEBS Journal (2006) Vol. 273 Supplement 1, 234-234

31st FEBS Congress Molecules in Health & Disease, Istanbul June 2006.

C Di Martino and **ML Pallotta** Mitochondria isolated from fresh and aged-dehydrated slices tubers of Jerusalem artichoke can synthesize NAD(P) from NMN externally added

Federazione Italiana Scienze della Vita, VIII National Congress October 2006

C Di Martino and **ML Pallotta** Proline accumulation in durum wheat seedlings under salt stress conditions depends on its own mitochondrial transport. Federazione Italiana Scienze della Vita, VIII National Congress October 2006

Pallotta ML The role of proline uptake in yeast mitochondria and the feast-famine regime.

Yeast, Vol.22 Issue S1 August 2005, Supplement " XXII International Conference on Yeast Genetics and Molecular Biology "Bratislava Slovak Republic.

Maria Luigia Pallotta, Daniela Valenti, Michelina Iacovino and Salvatore Passarella
Two separate pathways for D-lactate oxidation by *Saccharomyces cerevisiae* which differ in energy production and carrier involvement.
Biochim Biophys Acta -Bioenergetics. 2004;1608(2-3):104-13.

Pallotta ML Flavin adenine dinucleotide (FAD) and flavin mononucleotide (FMN) metabolism in *Saccharomyces cerevisiae* mitochondria.
Gene Transcription in Yeast: EuroConference. (European Science Foundation) Spain June 2004

Pallotta MLThe existence in *Saccharomyces cerevisiae* of a protective mechanism in response to methylglyoxal: the role of D-lactate movement across the inner mitochondrial membrane and its function on gluconeogenesis.
2004 Yeast Genetics & Molecular Biology University of Washington, Seattle, WA. July 27 - August 1, 2004.

ML. Pallotta, D. Valenti, M. Iacovino and S. Passarella D-lactate transport and metabolism in isolated *Saccharomyces cerevisiae* mitochondria.
"XXI International Conference on Yeast Genetics and Molecular Biology" Goteborg, Sweden. July 7-12, 2003. Yeast (2003) Jul, Supp 1, S11-390

Catello Di Martino, Roberto Pizzuto, **Maria Luigia Pallotta**, Donato Pastore, Aurelio De Santis and Salvatore Passarella Proline catabolism and mitochondrial transport: the occurrence of the proline uniporter and of the proline/glutamate antiporter in mitochondria isolated from durum wheat.
5° Convegno Federazione Italiana Scienze della Vita, 2003

Fратиани A., Pastore D., **Pallotta ML.**, Chiatante D. and Passarella S. Increase of membrane permeability of mitochondria isolated from water stress adapted potato cells.
Bioscience Reports (2001) 21: 81-91.

Pallotta ML., Fratianni A., Simone S. and Passarella S Fumarate/malate antiport in *Saccharomyces cerevisiae* mitochondria International Symposium on Molecular Basis of Biomembrane Monopoli (Ba) June 1999

Pallotta ML., Fratianni A. and Passarella S. Metabolites transport in isolated yeast mitochondria: fumarate/malate and succinate/malate antiports
FEBS Letters (1999) 462: 313-316

Fратиани A., **Pallotta ML.**, De Virgilio C., Chiatante D. and Passarella S. Water stress affects ADP/ATP and succinate/malate antiports in mitochondria isolated from potato (*Solanum tuberosum*) cell suspension cultures International Symposium on Molecular Basis of Biomembrane Monopoli (Ba) June 1999

Fратиани A., **ML. Pallotta**, De Virgilio C., Chiatante D. and Passarella S. The effect of water stress on certain features of ADP/ATP and Succinate/malate antiports in mitochondria isolated from *Solanum tuberosum* cell suspension cultures
GIBB, Scuola Superiore di Specializzazione Sant'Anna Pisa-September 1999.

ML. Pallotta, C. Brizio, A. Fratianni, C. De Virgilio, M. Barile, S. Passarella *Saccharomyces cerevisiae* mitochondria can synthesise FMN and FAD from externally added riboflavin and export them to the extramitochondrial phase
FEBS Letters (1998) 428: 245-249

A. Fratianni, **ML. Pallotta**, D. Chiatante, S. Passarella Effect of laser irradiation on the proliferative activity of *Phaseolus vulgaris* seedlings
Ital. J. Bioc. (1997) 46 (1) 1-60

D. Chiatante, A. Fratianni, **ML. Pallotta**, S. Passarella Increase in the proliferative activity of *Phaseolus vulgaris* seedlings by He-Ne Laser irradiation of embryos
Laser & Technology (1997) Vol. 7, N°1-2, 13-18.

ML. Pallotta, A. Fratianni, C. Brizio, C. De Virgilio, M. Barile e S. Passarella Processi di trasporto in mitocondri di *Saccharomyces cerevisiae*
GIBB-SIB Giugno 1997

ML. Pallotta, A. Fratianni D. Chiatante and S. Passarella
Effetto del laser ad Helium-Neon sulla crescita di piantule di *Phaseolus vulgaris*
II Convegno Nazionale Congiunto di Fotobiologia e Fotobiochimica Maratea Giugno 1996

A. Fratianni, **ML. Pallotta**, D. Chiatante S. Passarella
Alterazioni biochimiche in plantule di *Phaseolus vulgaris* derivate da embrioni irraggiati con laser ad Helium-Neon
Sezioni meridionali SIB Potenza 1996

G. Salzano, **ML. Pallotta**, M. F. Maddonni, R. Coppola, F. Villani, C. Sorlini Identification of *Listeria monocytogenes* in food and environmental by polymerase chain reaction
Journal of environmental science and health (1995) Part A: Environmental Science and Engineering & Toxic and Hazardous Substance Control A 30 (1): 63-71

F. Pilla, S. Dell'Aquila, L. Taibi, C. Tripaldi, S. Puppo, F. Napolitano, **ML. Pallotta**, M. Angelucci, A. Girolami Influenza del polimorfismo genetico della Beta-lattoglobulina su alcune caratteristiche chimico-fisiche e tecnologiche del latte di pecora
XI Congresso A. S. P. A. Grado Giugno 1995.

S. Torriani, **ML. Pallotta**
Use of Polymerase Chain Reaction to detect *Listeria monocytogenes* in silages.
Biotechnology Techniques (1994) Vol 8 N° 3: 157-160

ML. Pallotta, A. Bellitti, F. Pilla e A. Girolami
Distribuzione degli alleli A e B mediante PCR (Polymerase Chain Reaction) e RFLP (Restriction fragment length polymorphism).

. Biotechnology for milk production Torino Novembre 1994

G. Salzano, R. Coppola, M.F. Maddonni, **ML. Pallotta**, F. Villani e C. Sorlini.
Identification of *Listeria monocytogenes* in Mozzarella Cheese by Polymerase Chain Reaction
employment.
Ann. Microbiol. Enzimol. (1993) 43: 159-163.

G. Salzano, R. Coppola, M.F. Maddonni, **ML. Pallotta**, F. Villani e C. Sorlini Detection of *Listeria*
monocytogenes in environment and food by means of Polymerase Chain Reaction in technique.
Sixth European Congress on Biotechnology Firenze 1993.

Dellaglio, S. Torriani, F. Gardini, **ML. Pallotta**.

Use of Polymerase Chain Reaction" (PCR) to detect *Listeria monocytogenes* in silage and
modelling of the growth in relation of some chemico-physical factors.

Silage Reserch 1993. 10th International Conference on Silage Research. Dublin (Ireland)

F. Dellaglio, V. Giaccone, S. Torriani, G. Salzano, G. Colavita, A. Mazzeo and **ML. Pallotta** The
Polymerase Chain Reaction (PCR) as a rapid method to detect *Salmonella* spp. in fresh meat and
meat products. The International Committee on Food Microbiology and Hygiene Bingen (Germany)
1993.

F. Dellaglio, V. Giaccone, S. Torriani, G. Colavita, G. Salzano and **ML. Pallotta**.

The Polymerase Chain Reaction (PCR) as a rapid method to detect *Salmonella* spp. in red meat.
The International Committee on Food Microbiology and Hygiene Bingen (Germany) 1993.

S. Torriani, G. Salzano, **ML. Pallotta** e F. Dellaglio Impiego della "Polymerase Chain Reaction"
(PCR) per rilevare *Listeria monocytogenes* negli insilati.

S. I. M. 24° Congresso Nazionale. Genova 1992.

G. Salzano, **ML. Pallotta**, M. F. Maddonni, F. Villani, R. Coppola e C. Sorlini Uso della
"Polymerase Chain Reaction" (PCR) per rilevare *Listeria monocytogenes* negli alimenti.

S. I. M . 24° Congresso Nazionale. Genova 1992.

Chapter in book

Bacteriocins: Production, Applications and Safety

Edited by Troy Padilla

Chapter 1 : Bacteriocin Producing Cultures: A Sustainable Way for Food Safety Improvement and Probiotics with Additional Health Promoting Effects (2016) ISBN: 978-1-63484-531-1

F Rossi, **ML Pallotta**